



Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Source

Public Health outcomes framework

NHS outcomes framework

Adult social care outcomes framework

National average

Dudley LA peers

Worst

Dudley LA peers

Best

25th Pecentile 75th

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|-------------|-------------------|--|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 0.1i males | Healthy male life expectancy at birth 2010-12 (Years) | | 62.8 | 63.4 | 52.5 | | 70.0 | • |
| | 0.1i females | Healthy female life expectancy at birth 2010-12 (Years) | | 64.7 | 64.1 | 55.5 | ♦ | 71.0 | |
| | 0.1ii males | Male life expectancy at birth 2010 - 12 (Years) | | 78.9 | 79.2 | 74.0 | | 82.1 | |
| | 0.1ii females | Female life expectancy at birth 2010 - 12 (Years) | | 83.0 | 83.0 | 79.5 | • | 85.9 | • |
| | 0.2ii Females | Number of upper tier local authorities for which the local slope index of inequality in female life expectancy (as defined in 0.2iii) has decreased | | | 73.0 | | | | • |
| Đ. | 0.2ii Males | Number of upper tier local authorities for which the local slope index of inequality in male life expectancy (as defined in 0.2iii) has decreased (Number) | | | 73.0 | | | | • |
| Overarching | 0.2iii males | Slope index of inequality in male life expectancy at birth within English local authorities, based on local deprivation deciles within each area 2010 - 12 | | 9.5 | | 3.9 | | 16.0 | • |
| Ove | 0.2iii females | Slope index of inequality in female life expectancy at birth within English local authorities, based on local deprivation deciles within each area 2010 - 12 | | 5.8 | | 1.3 | 0 | 11.4 | |
| | 1bi | Life expectancy at 75, males 2010-12 (Years) | | 11.4 | 11.5 | 10.0 | ∞ | 13.6 | • |
| | 1bii | Life expectancy at 75, females 2010-12 (Years) | | 13.3 | 13.3 | 11.6 | | 15.9 | • |
| | 4.3 Males | Mortality rate from causes considered preventable (Males) 2010-12 (Rate per 100,000 population) | 1096 | 250.3 | 238.4 | 430.9 | • | 164.9 | |
| | 4.3 Females | Mortality rate from causes considered preventable (Females) 2010-12 (Rate per 100,000 population) | 621 | 133.7 | 140.6 | 253.9 | | 94.7 | |
| | 4.3 Persons | Mortality rate from causes considered preventable (Persons) 2010-12 (Rate per 100,000 population) | 1717 | 190.3 | 187.8 | 340.5 | | 136.2 | • |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Source

Public Health outcomes framework

NHS outcomes framework

Adult social care outcomes framework

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|-------------|------------------|--|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 4.4i Males | Mortality rate from all cardiovascular diseases (including heart diseases and stroke) in Males less than 75 years of age 2010-12 (Rate per 100,000 | 460 | 114.0 | 114.0 | 204.4 | • | 79.3 | |
| | 4.4i Females | Mortality rate from all cardiovascular diseases (including heart diseases and stroke) in Females less than 75 years of age 2010-12 (Rate per 100,000 population) | 198 | 47.5 | 50.1 | 88.4 | ** | 28.6 | |
| | 4.4i Persons | Mortality rate from all cardiovascular diseases (including heart diseases and stroke) in Persons less than 75 years of age 2010-12 (Rate per 100,000 | 658 | 80.3 | 81.1 | 144.7 | • | 55.7 | |
| | 1.1 Persons | Under 75 mortality rate from cardiovascular disease 2012 (Rate per 100,000 population) | | 74.2 | 77.7 | 130.0 | ◇ | 35.7 | |
| | 4.4ii Males | Mortality rate from all cardiovascular diseases (including heart disease and stroke) that is considered preventable in Males less than 75 years of age 2010- | 443 | 54.1 | 53.5 | 95.2 | ♦ | 29.3 | |
| Overarching | 4.4ii Females | Mortality rate from all cardiovascular diseases (including heart disease and stroke) that is considered preventable in Females less than 75 years of age | 102 | 24.7 | 27.6 | 54.4 | → | 15.0 | |
| Overa | 4.4ii Persons | Mortality rate from all cardiovascular diseases (including heart disease and stroke) that is considered preventable in Persons less than 75 years of age 2010-12 (Rate per 100 000 population) | 443 | 54.1 | 53.5 | 95.2 | • | 29.3 | • |
| | 4.5i Males | Mortality rate from all cancers for Males less than 75 years of age 2010-12 | 713 | 176.5 | 163.6 | 238.9 | • | 122.8 | |
| | 4.5i Females | Mortality rate from all cancers for Females less than 75 years of age 2010-12 (Rate per 100,000 population) | 573 | 136.4 | 130.8 | 181.3 | • | 105.3 | |
| | 4.5i Persons | Mortality rate from all cancers for Persons less than 75 years of age 2010-12 (Rate per 100,000 population) | 1286 | 155.9 | 146.5 | 207.3 | • | 113.5 | • |
| | 4.5ii Males | Mortality rate from all cancers that is considered preventable in Males less than 75 years of age 2010-12 (Rate per 100,000 population) | 405 | 100.1 | 92.7 | 154.4 | • | 53.1 | |
| | 4.5ii Females | Mortality rate from all cancers that is considered preventable in Females less than 75 years of age 2010-12 (Rate per 100,000 population) | 320 | 76.0 | 77.9 | 121.4 | • | 54.6 | |



Domain Indicator

4.5ii Persons

4.6i Males

4.6i Females

4.6i Persons

1.3

Persons

4.6ii

Males

4.6ii **Females**

4.6ii Persons

4.7i Males

4.7i Females

4.7i

Persons

1.2

Overarching



HEALTH AND WELLBEING PRIORITIES -OUTCOMES FRAMEWORKS

Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be

Public Health outcomes framework

NHS outcomes framework

Source

| ic | Adult social care outcomes framework Adult social care outcomes framework | | | | | | | | | | | |
|----|---|--------------------|----------------|------------|--------------|---------------|----------|----------------|--|--|--|--|
| or | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value | | | | |
| 6 | Mortality rate from all cancers that is considered preventable in Persons less than 75 years of age 2010-12 (Rate per 100,000 population) | 725 | 87.7 | 84.9 | 134.9 | • | 53.8 | | | | | |
| es | Mortality rate from liver disease in Males less than 75 years of age 2010-12 (Rate per 100,000 population) | 131 | 31.8 | 23.7 | 58.4 | • • | 13.0 | | | | | |
| S | Mortality rate from liver disease in Females less than 75 years of age 2010-12 (Rate per 100,000 population) | 77 | 18.8 | 12.6 | 25.0 | • • | 6.9 | | | | | |
| 6 | Mortality rate from liver disease in Persons less than 75 years of age 2010-12 (Rate per 100,000 population) | 208 | 25.2 | 18.0 | 41.6 | | 10.3 | • | | | | |
| 6 | Under 75 mortality rate from liver disease 2012 (Rate per 100,000 population) | | 27.4 | 17.7 | 40.9 | • • | 3.1 | • | | | | |
| | Mortality rate from liver disease that is considered preventable in Males less than 75 years of age 2010-12 (Rate per 100,000 population) | 121 | 29.3 | 21.1 | 54.9 | • • | 10.8 | | | | | |
| 3 | Mortality rate from liver disease that is considered preventable in Females less than 75 years of age 2010-12 (Rate per 100,000 population) | 69 | 16.8 | 10.6 | 21.4 | • • | 6.3 | | | | | |
| 6 | Mortality rate from liver disease that is considered preventable in Persons less than 75 years of age 2010-12 (Rate per 100,000 population) | 190 | 23.0 | 15.8 | 38.2 | | 9.0 | • | | | | |
| s | Mortality rate from respiratory diseases in Males less than 75 years of age 2010-12 (Rate per 100,000 population) | 164 | 40.8 | 39.6 | 92.1 | | 24.6 | | | | | |
| 3 | Mortality rate from respiratory diseases in Females less than 75 years of age 2010-12 (Rate per 100,000 population) | 119 | 28.4 | 27.9 | 71.5 | | 12.6 | | | | | |
| 6 | Mortality rate from respiratory diseases in Persons less than 75 years of age 2010-12 (Rate per 100,000 population) | 283 | 34.5 | 33.5 | 81.6 | | 20.5 | | | | | |
| 3 | Under 75 mortality rate from respiratory disease 2012 (Rate per 100,000 population) | | 29.3 | 32.7 | 81.6 | 0 | 7.2 | • | | | | |



Domain Indicator

4.7ii

Males

4.7ii

Females

4.7ii

Persons

4.8 -Males

4.8 -

Females

4.8 -Persons

4.14i

4.14ii

4.14iii

4.15i

3.5ii

1a males

Overarching



HEALTH AND WELLBEING PRIORITIES -OUTCOMES FRAMEWORKS

Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

population)

population)

population)

No significance can be calculated

Source
Public Health outcomes framework

NHS outcomes framework

Mortality rate from respiratory diseases that is considered preventable in Males

Mortality rate from respiratory diseases that is considered preventable in

Females less than 75 years of age 2010-12 (Rate per 100,000 population)

Mortality rate from respiratory diseases that is considered preventable in

Persons less than 75 years of age 2010-12 (Rate per 100,000 population)

Mortality rate from communicable diseases in Males (Rate per 100,000

Mortality rate from communicable diseases in Persons (Rate per 100,000

Mortality rate from communicable diseases in Females (Rate per 100,000

Emergency admissions for fractured neck of femur in persons aged 65 and

Emergency admissions for fractured neck of femur in persons aged 65-79

Emergency admissions for fractured neck of femur in persons aged 80 and

Potential Years of Life Lost from causes considered amenable to healthcare

Excess winter deaths single year all ages Aug 11 - Jul 12 (Ratio)

Incidence of Tuberculosis 2010-12 (Rate per 100,000 population)

over 2012/13 (Rate per 100,000 population)

over 2012/13 (Rate per 100,000 population)

(males) 2012 (Rate per 100,000 population)

2012/13 (Rate per 100,000 population)

less than 75 years of age 2010-12 (Rate per 100,000 population)

Adult social care outcomes framework

Local

Numerator

78

58

136

370

116

2490.6 2556.2

4581.8

National average

Dudley LA peers

Worst

Best

| | | | 25th Pecentile 75th | | |
|---------------|-----------|--------------|---------------------|----------|----------------|
| Loca Value | | Eng Worst | England Range | Eng Best | Previous value |
| 19 | .3 20.1 | 50.4 | • | 10.6 | |
| 13 | .9 15.2 | 40.2 | | 7.3 | |
| 16 | 17.6 | 45.0 | | 7.9 | • |
| | | | | | |
| | | | | | |
| | | | | | • |
| 596 | 5.6 568.1 | 808.4 | • | 403.1 | • |
| 249 | .4 237.3 | 401.7 | | 121.8 | |
| 1603 | .2 1527.6 | 2150.0 | • | 1107.7 | |
| 12 | 16.1 | 30.7 | ∞ | 2.1 | |
| 10 | .5 15.1 | 112.3 | 0 | 0.0 | • |
| | | | | | |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------|---------------|--|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 1a females | Potential Years of Life Lost from causes considered amenable to healthcare (females) 2012 (Rate per 100,000 population) | | 2122.1 | 2063.2 | 3368.2 | | 547.8 | • |
| | 1.4i | One-year survival from all cancers (%) | | | 68.2 | | | | |
| Overarching | 1.4ii | Five-year survival from all cancers (%) | | | 47.9 | | | | |
| Overa | 1.4iii | One-year survival from colorectal cancer (%) | | | 69.5 | | | | |
| | 1.4iv | Five-year survival from colorectal cancer (%) | | | 51.4 | | | | |
| | 1.7 | Premature mortality for people with a learning disability (TBC) | | | | | | | |
| | 1.1i | Children in poverty (all dependent children under 20) 2011 (%) | 15005 | 22.1 | 20.1 | 46.1 | | 2.9 | • |
| Ç | 1.1ii | Children in poverty (under 16s) 2011 (%) | 13095 | 22.8 | 20.6 | 43.6 | • | 2.8 | |
| Childre | 1.2ia | School Readiness: The percentage of children achieving a good level of development at the end of reception 2012/13 (%) | 1928 | 51.1 | 51.7 | 27.7 | • | 69.0 | • |
| Healthy Children | 1.2ib | School Readiness: The percentage of children with free school meal status achieving a good level of development at the end of reception 2012/13 (%) | 261 | 33.9 | 36.2 | 17.8 | • | 60.0 | |
| Ĭ | 1.2iia | School Readiness: The percentage of Year 1 pupils achieving the expected level in the phonics screening check 2012/13 (%) | 2467 | 68.0 | 69.1 | 58.8 | ○ ♦ | 79.0 | |
| | 1.2iib | School Readiness: The percentage of Year 1 pupils with free school meal status achieving the expected level in the phonics screening check 2012/13 (%) | 400 | 52.4 | 55.8 | 37.2 | • • | 70.9 | |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

<u>Source</u>

Public Health outcomes framework

NHS outcomes framework

Adult social care outcomes framework

National average

Dudley LA peers

Worst

Best

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------|-----------|---|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 1.3 | Pupil absence 2012-13 (%) | | 5.6 | 5.3 | 6.3 | • | 3.6 | • |
| | 1.4 | First time entrants to the youth justice system 2013 (Rate per 100,000 population) | | 384.5 | 440.9 | 846.5 | (| 171.0 | • |
| | 1.5 | 16-18 year olds not in education employment or training 2013 (%) | | 5.9 | 5.3 | 9.8 | • | 1.8 | • |
| | 2.8 | Emotional well-being of looked after children 2012/13 (Score out of 40) | | 12.8 | 14.0 | 9.4 | • | 21.5 | • |
| C | 2.1 | Low birth weight of term babies 2011 (%) | | 3.4 | 2.8 | 5.3 | • | 1.6 | • |
| Childre | 4.1 | Infant mortality 2010 - 12 (Rate per 1,000 live births) | 50 | 4.4 | 4.1 | 7.5 | ♦ | 0.0 | • |
| Healthy Children | 1.6i | Infant mortality 2012 (Rate per 1,000 live births) | 19 | 4.8 | 4.0 | 10.5 | • | 0.0 | • |
| Ĭ | 1.6iii | Five year survival from all cancers in children (TBC) | | | 81.3 | | | | |
| | 5.5 | Admission of full term babies to neonatal care 2011 (%) | 165 | 4.8 | 6.0 | 25.1 | • | 0.9 | • |
| | 3.2 | Emergency admissions for children with Lower Respiratory Tract Infections (LRTIs) 2012-13 (Rate per 100,000 population) | | 584.1 | 371.2 | 745.0 | • | 115.9 | • |
| | 5.6 | Incidence of harm to children due to 'failure to monitor' (Number) | | | 1017.0 | | | | |
| | 4.8 | Children and young people's experience of healthcare (TBC) | | | | | | | |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Source
Public Health outcomes framework

NHS outcomes framework

Adult social care outcomes framework

National average

Dudley LA peers

Worst ● ◆ Best

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------|-----------------|--|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 2.6i | Excess weight in 4-5 and 10-10 year olds - 4-5 year olds 2012-13 (%) | 865 | 23.6 | 22.2 | 32.2 | • | 16.1 | • |
| | 2.6ii | Excess weight in 4-5 and 10-11 year olds - 10 -11 year olds 2012-13 (%) | 1244 | 37.1 | 33.3 | 44.2 | • ♦ | 24.1 | • |
| | 4.2 | Tooth decay in children aged 5 2011-12 (%) | | 0.6 | 0.9 | 2.1 | ♦ ○ | 0.0 | • |
| | 2.7ia | Hospital admissions caused by unintentional and deliberate injuries in children (aged 0-14 years) 2012-13 (Rate per 10,000 population) | 525 | 94.9 | 103.8 | 191.3 | •• | 61.7 | |
| C. | 2.7ib | Hospital admissions caused by unintentional and deliberate injuries in children (aged 0-4 years) 2012/13 () | 259 | 135.6 | 134.7 | 282.4 | • | 76.0 | |
| Healthy Children | | Hospital admissions caused by unintentional and deliberate injuries in young people (aged 15-24) 2012-13 (Rate per 10,000 population) | 433 | 115.6 | 130.7 | 277.3 | \Q | 63.8 | |
| ealthy | 3.2ii Males | Chlamydia diagnoses Males - CTAD Data 2013 (Rate per 100,000 population) | 212 | 1109.6 | 1387.5 | 4262.0 | © | 599.4 | |
| Ĭ | | Chlamydia diagnoses Females - CTAD Data 2013 (Rate per 100,000 population) | 456 | 2482.8 | 2633.5 | 1093.7 | • | 6358.2 | |
| | | Chlamydia diagnoses Persons - CTAD Data 2013 (Rate per 100,000 population) | 672 | 1793.3 | 2015.6 | 5758.5 | • | 840.0 | • |
| | 2.4i | Under 18 conceptions 2012 (Rate per 1,000 female population aged 15-17) | 203 | 34.6 | 27.7 | 52.0 | •• | 14.2 | • |
| | 2.4ii | Under 18 conceptions: conceptions in those aged under 16 2011 (Rate per 1,000 female population aged 13-15) | | 6.7 | 6.1 | 2.2 | | 11.5 | |
| | 3.3i (1yr olds) | Hepatitis B vaccination coverage (one year olds) 2012-13 (%) | 4 | 100.0 | _ | 40.0 | | 100.0 | |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Source
Public Health outcomes framework

NHS outcomes framework

Adult social care outcomes framework

National average

Dudley LA peers

Worst

Best

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------|-------------------------|---|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 3.3i (2yr olds) | Hepatitis B vaccination coverage (two year olds) 2012-13 (%) | 4 | 100.0 | | 9.1 | | 100.0 | |
| | 3.3ii | BCG vaccination coverage (aged under 1 year) 2011-12 (%) | 518 | 100.0 | 96.5 | 2.1 | | 100.0 | • |
| | 3.3iii (1 year olds) | DTaP/IPV/Hib vaccination coverage (one year old) 2012-13 (%) | 3640 | 97.4 | 94.7 | 79.0 | • | 99.0 | |
| | 3.3iii (2 year olds) | DTaP/IPV/Hib vaccination coverage (two year old) 2012-13 (%) | 3569 | 97.9 | 96.3 | 81.9 | ∞ | 99.4 | |
| L. | 3.3iv | MenC vaccination coverage (one year olds) 2012-13 (%) | 3629 | 97.1 | 93.9 | 75.9 | > • | 98.8 | • |
| Healthy Children | 3.3v | PCV vaccination coverage (one year olds) 2012-13 (%) | 3645 | 97.5 | 94.4 | 78.7 | • • | 99.0 | • |
| ealthy | 3.3vi (2 Year olds) | Hib/MenC booster vaccination coverage (two year olds) 2012-13 (%) | 3467 | 95.1 | 92.7 | 77.0 | < o | 98.3 | |
| Ĭ | 3.3vi (5 Year olds) | Hib/MenC booster vaccination coverage (five year olds) 2012-13 (%) | 3363 | 93.4 | 91.5 | 75.7 | | 98.1 | |
| | 3.3vii | PCV booster vaccination coverage (two year olds) 2012-13 (%) | 3522 | 96.7 | 92.5 | 75.1 | ⋄ | 97.5 | • |
| | 3.3viii | MMR vaccination coverage for one dose (two year olds) 2012-13 (%) | 3502 | 96.1 | 92.3 | 77.4 | ⋄ | 98.4 | • |
| | 3.3ix | MMR vaccination coverage for one dose (five year olds) 2012-13 (%) | 3427 | 95.2 | 93.9 | 82.1 | • | 98.3 | |
| | 3.3x | MMR vaccination coverage for two doses (five year olds) 2012-13 (%) | 3207 | 89.1 | 87.7 | 68.9 | O | 97.0 | • |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Source
Public Health outcomes framework

NHS outcomes framework

Adult social care outcomes framework

National average

Dudley LA peers

Worst ● ◆ Best

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|---------------------|-----------|---|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| Healthy Children | 3.3xi | Td/IPV booster vaccination coverage (13-18 year olds) 2011-12 (%) | 2893 | 12.1 | 11.4 | 2.7 | ♦ ⊃ | 24.7 | • |
| Hea | 3.3xii | HPV vaccination coverage (females 12-13 year olds) 2012-13 (%) | 1617 | 92.0 | 86.1 | 62.1 | ◇ | 96.2 | • |
| | 2.2i | Breastfeeding - Breastfeeding initiation 2012-13 (%) | 2028 | 54.1 | 73.9 | 40.8 | • • | 94.7 | • |
| | 2.2ii | Breastfeeding - Breastfeeding prevalence at 6-8 weeks after birth 2012-13 (%) | 1073 | 28.0 | 47.2 | 17.5 | • • | 83.3 | • |
| | 2.3 | Smoking status at time of delivery 2012-13 (%) | 535 | 14.3 | 12.7 | 30.8 | • | 2.3 | • |
| es | 2.14i | Smoking Prevalence 2012 (%) | | 18.7 | 19.5 | 29.8 | • | 12.1 | |
| Healthy Lifestyles | 2.14ii | Smoking prevalence - routine & manual 2012 (%) | | 27.7 | 29.7 | 44.3 | | 14.2 | |
| ealthy I | 2.12 | Excess weight in adults 2012 (%) | 565 | 70.2 | 63.8 | 74.4 | • • | 45.9 | • |
| <u>¥</u> | 2.13i | Percentage of physically active and inactive adults - active adults 2013 (%) | | 46.5 | 55.6 | 43.4 | • • | 71.2 | • |
| | 2.13ii | Percentage of active and inactive adults - inactive adults 2013 (%) | | 36.7 | 28.9 | 39.2 | • • | 16.3 | • |
| | 2.18 | Alcohol related admissions to hospital 2012/13 (Rate per 100,000 population) | 2381 | 777.7 | 636.9 | 1120.6 | • • | 365.0 | • |
| | 2.20i | Cancer screening coverage - breast cancer 2013 (%) | 25493 | 75.1 | 76.3 | 58.2 | | 84.5 | • |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|--------------------|-----------|---|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 2.20ii | Cancer screening coverage - cervical cancer 2013 (%) | 56548 | 73.8 | 73.9 | 58.6 | • | 83.2 | • |
| styles | 2.21vii | Access to non- cancer screening programmes diabetic retinopathy 2011-12 (%) | 10320 | 72.9 | 80.9 | 66.7 | • • | 95.0 | • |
| Healthy Lifestyles | 2.22iii | Take up of NHS Health Check Programme by those eligible - health check offered 2013/14 (%) | 19903 | 20.9 | 18.5 | 0.8 | | 44.4 | • |
| Health | 2.22iv | Take up of NHS Health Check programme by those eligible - health check take up of those offered 2013/14 (%) | 7867 | 39.5 | 49.0 | 14.6 | •• | 100.0 | • |
| | 2.22v | Take up of NHS Health Check programme by those eligible - health check take up 2013/14 (%) | 7867 | 8.2 | 9.0 | 0.9 | • | 29.1 | |
| | 2.10 | Self-harm 2011-12 (Rate per 100,000 population) | 684 | 234.3 | 207.9 | 542.4 | • | 51.2 | • |
| | 4.9 | Excess under 75 mortality in adults with serious mental illness 2011/12 (Rate per 100,000 population) | | 342.8 | 337.4 | 510.4 | | 124.7 | • |
| spu | 2.23i | Self-reported well- being - people with a low satisfaction score 2012-13 (%) | | 4.2 | 5.8 | 10.1 | ♦ • | 3.4 | • |
| Healthy Minds | 2.23ii | Self-reported well- being - people with a low worthwhile score 2012-13 (%) | | 4.6 | 4.4 | 8.2 | • | 2.9 | • |
| Hea | 2.23iii | Self-reported well- being - people with a low happiness score 2012-13 (%) | | 8.9 | 10.4 | 15.8 | • | 5.5 | • |
| | 2.23iv | Self-reported well- being - people with a high anxiety score 2012-13 (%) | | 19.6 | 21.0 | 29.0 | • | 10.9 | |
| | 2 | Health related quality of life for people with long-term conditions 2012/13 (Number) | 1917 | 0.7 | 0.7 | 0.6 | • ◊ | 0.8 | • |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

National average

Dudley LA peers

Worst

Best

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England | Range | Eng Best | Previous value |
|---------------|--------------|---|--------------------|----------------|------------|--------------|-----------|-------------|----------|----------------|
| | 2.4 | Health-related quality of life for carers 2012-13 (Number) | 910 | 0.8 | 0.8 | 0.8 | | | 0.9 | • |
| | | Percentage of male adults in contact with secondary mental health services who live in stable and appropriate accommodation 2011/12 (%) | 500 | 50.8 | 53.8 | 3.1 | • | | 91.1 | |
| | | Percentage of female adults in contact with secondary mental health services who live in stable and appropriate accommodation 2011/12 (%) | 465 | 46.2 | 55.5 | 2.7 | • | | 94.2 | |
| | | Percentage of adults in contact with secondary mental health services who live in stable and appropriate accommodation 2012/13 (%) | 600 | 30.0 | 58.5 | 5.5 | | > | 94.1 | • |
| | 1.8iii | Gap in the employment rate for those in contact with secondary mental health services and the overall employment rate 2012/13 (%) | | 69.2 | 62.3 | 75.1 | | ♦ | 53.1 | |
| Healthy Minds | 11 1 | Proportion of adults in contact with secondary mental health services in paid employment 2013-14 (%) | | 4.4 | 7.1 | 1.6 | | ♦ | 17.5 | • |
| Healthy | 1 61 1/12126 | Male adults with a learning disability who live in stable and appropriate accommodation 2012/13 (%) | 300 | 70.7 | 73.2 | 32.2 | \$ | | 97.9 | |
| | | Female adults with a learning disability who live in stable and appropriate accommodation 2012/13 (%) | 190 | 68.8 | 74.0 | 33.3 | • | | 97.9 | |
| | | Adults with a learning disability who live in stable and appropriate accommodation 2012/13 (%) | 495 | 70.0 | 73.5 | 32.6 | ◇● | | 96.6 | • |
| | | Proportion of adults in contact with secondary mental health services living independently, with or without support 2013-14 (%) | | 52.4 | 60.9 | 12.6 | • | ♦ | 93.3 | • |
| | 2.6i | Estimated diagnosis rate for people with dementia (%) | | | 48.7 | | | | | |
| | 2 bil | Effectiveness of dementia post-diagnosis care in sustaining independence and improving quality of life (TBC) | | | | | | | | |





HEALTH AND WELLBEING PRIORITIES -OUTCOMES FRAMEWORKS

Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

| Domain | Indicator | · | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------------|-----------|--|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 4.7 | Patient experience of community mental health services 2013 (Score out of 100) | | 87.6 | 85.8 | 80.9 | • | 90.9 | • |
| | 1.10 | Killed and seriously injured casualties on England's roads 2010-12 (Rate per 100,000 population) | 283 | 30.1 | 40.5 | 665.6 | | 0.0 | • |
| | 3.1 | Fraction of mortality attributable to particulate air pollution 2012 (%) | | 5.2 | 5.1 | 8.3 | | 3.0 | • |
| | 1.16 | Utilisation of outdoor space for exercise/health reasons Mar 2012- Feb 2013 (%) | | 14.3 | 15.3 | 0.0 | • | 41.2 | • |
| spoor | 3.6 | NHS organisations with a board approved sustainable development management plan 2012/13 (%) | 2 | 40.0 | 59.0 | 16.7 | • • | 100.0 | • |
| Healthy Neighbourhoods | 1.12ii | Violent crime (including sexual violence) - violence offences per 1,000 population 2012-13 (Rate per 1,000 population) | 2157 | 6.9 | 10.6 | 69.9 | • | 4.1 | • |
| y Neig | 1.12i | Violent crime (including sexual violence) - hospital admissions for violence 2010-11/2012-13 (Rate per 100,000 population) | 583 | 62.2 | 57.6 | 167.8 | | 9.3 | • |
| Health | 1.12iii | Violent crime (including sexual violence) - Rate of sexual offences per 1,000 population 2012-13 (%) | 262 | 0.8 | 0.8 | 4.7 | • | 0.3 | |
| | 1.13i | Re-offending levels percentage of offenders who re 2011 (%) | 548 | 23.1 | 26.9 | 36.3 | ♦ • | 14.4 | • |
| | 1.13ii | Re-offending levels average number of re 2011 (Mean average) | 1554 | 0.7 | 0.8 | 1.3 | ♦ • | 0.3 | • |
| | 1.14i | The percentage of the population affected by noise - Number of complaints about noise 2011-12 (Rate per 1,000 population) | 1528 | 4.9 | 7.5 | 116.0 | | 1.4 | |
| | 1.14ii | The percentage of the population exposed to road, rail and air transport noise of 65dB(A) or more, during the daytime 2011 (%) | 16410 | 5.2 | 5.2 | 28.5 | • | 0.0 | |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------------|------------------|--|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| Healthy Neighbourhoods | 1.14iii | The percentage of the population exposed to road, rail and air transport noise of 55 dB(A) or more during the night- time 2011 (%) | 22020 | 7.0 | 8.0 | 42.4 | | 0.0 | |
| | 1.15i | Statutory homelessness - homelessness acceptances 2012/13 (Rate per 1,000 households) | 171 | 1.3 | 2.4 | 11.4 | ♦ • | 0.2 | • |
| | 1.15ii | Statutory homelessness - households in temporary accommodation 2012/13 (Rate per 1,000 households) | 55 | 0.4 | 2.4 | 33.2 | | 0.0 | • |
| Health | 1.17 | Fuel Poverty 2012 (%) | 18129 | 14.3 | 10.4 | 21.3 | ◆● | 2.4 | • |
| | 2.17 | Recorded diabetes 2012-13 (%) | 16515 | 6.5 | 6.0 | 8.4 | ◇● | 3.7 | • |
| | 4.11 Males | Emergency readmissions within 30 days of discharge from hospital (Male) 2011/12 () | 2107 | 12.0 | 12.1 | 14.9 | | 8.7 | |
| Se | 4.11 Females | Emergency readmissions within 30 days of discharge from hospital (Female) 2011/12 (%) | 2180 | 11.6 | 11.5 | 14.7 | | 8.3 | |
| Services | 4.11 Persons | Emergency readmissions within 30 days of discharge from hospital (Persons) 2011/12 () | 4287 | 11.8 | 11.8 | 14.5 | | 7.9 | |
| Healthy | 2.15i | Successful completion of drug treatment - opiate users 2012 (%) | 89 | 9.2 | 8.2 | 0.0 | ◇ • | 17.6 | • |
| Ĭ | 2.15ii | Successful completion of drug treatment - non-opiate users 2012 (%) | 80 | 44.0 | 40.2 | 0.0 | • | 68.4 | • |
| | 2.24i Females | Injuries due to falls in people aged 65 and over (Females) 2012/13 (Rate per 100,000 population) | 1050 | 2981.7 | 2419.9 | 4041.3 | • | 1452.1 | |
| | 2.24i Males | Injuries due to falls in people aged 65 and over (Males) 2012/13 (Rate per 100,000 population) | 481 | 2009.6 | 1602.1 | 2974.8 | • | 902.9 | |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

National average

Dudley LA peers

Worst

Dudley LA peers

Best

25th Pecentile 75th

| Domain | Indicator | • | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|----------|------------------|--|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 2.24i Persons | Injuries due to falls in people aged 65 and over (Persons) 2012/13 (Rate per 100,000 population) | 1531 | 2495.6 | 2011.0 | 3508.0 | • | 1177.5 | • |
| | 2.24ii | Emergency hospital admissions for injuries due to falls in persons aged 65 to 79 2012/13 (Rate per 100,000 population) | 525 | 1194.9 | 975.0 | 1826.4 | • • | 544.3 | • |
| | 2.24iii | Emergency hospital admissions for injuries due to falls in persons aged 80 and over 2012/13 (Rate per 100,000 population) | 1006 | 6267.9 | 5015.3 | 9118.7 | • | 2875.6 | • |
| | 2.3i | Unplanned hospitalisation for chronic ambulatory care sensitive conditions 2012-13 (Rate per 100,000 population) | | 855.8 | 820.5 | 1596.7 | | 217.3 | • |
| S | 2.3ii | Unplanned hospitalisation for asthma, diabetes and epilepsy in under 19s 2012-13 (Rate per 100,000 population) | | 341.3 | 340.6 | 892.8 | ♦ | 0.0 | • |
| Services | 3a | Emergency admissions for acute conditions that should not usually require hospital admission 2012-13 (Rate per 100,000 population) | | 1612.5 | 1204.3 | 2209.1 | • | 299.7 | • |
| Healthy | 3b | Emergency readmissions within 30 days of discharge from hospital 2011-12 (%) | | 11.8 | | 14.5 | | 7.9 | |
| Ĭ | 3.1i | Total health gain as assessed by patients for elective procedures - Unilateral Hip Replacements 2011-12 (Health score) | | 0.4 | 0.4 | 0.3 | | 0.5 | • |
| | 3.1ii | Total health gain as assessed by patients for elective procedures - Unilateral Knee Replacements 2011-12 (Health score) | | 0.3 | 0.3 | 0.2 | • | 0.4 | • |
| | 3.1iii | Total health gain as assessed by patients for elective procedures - Groin Hernia Surgery 2011-12 (Health score) | | 0.0 | 0.1 | 0.0 | • | 0.1 | • |
| | 3.1iv | Total health gain as assessed by patients for elective procedures - Varicose Vein Surgery 2011-12 (Health score) | | 0.1 | 0.1 | 0.1 | ♦ • | 0.2 | • |
| | 3.1v | Total health gain as assessed by patients for elective procedures - Psychological Therapies (Health score) | | _ | _ | _ | | | |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Source
Public Health outcomes framework

NHS outcomes framework

Adult social care outcomes framework

National average

Dudley LA peers

Worst ◆ Best

| Domain | Indicator | • | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|----------|-----------|---|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 3.3 | Proportion of people who recover from major trauma (Adjusted odds) | | | | | | | |
| | 3.4 | Proportion of stroke patients reporting an improvement in activity/lifestyle on the Modified Rankin Scale at 6 months (Score out of 6) | | | | | | | |
| | 3.5i | The proportion of patients with a fragility fracture recovering to their previous levels of mobility at 30 days (%) | | | 21.7 | | | | |
| | 3.5ii | The proportion of patients with a fragility fracture recovering to their previous levels of mobility at 120 days (%) | | | 47.3 | | | | |
| Se | 3.6i | The proportion of older people (65 and over) who were still at home 91 days after discharge from hospital into rehabilitation 2012-13 (%) | 220 | 87.4 | 81.4 | 53.7 | ♦ | 98.1 | • |
| Services | 3.6ii | The proportion of older people (65 and over) who were still at home 91 days after discharge from acute or community hospital into reablement/rehabilitation services offered rehabilitation 2012-13 (%) | 255 | 2.8 | 3.2 | 0.4 | • | 25.4 | |
| Healthy | 4ai | Patient experience of primary care - GP services 2012-13 (Score out of 100) | 4663 | 85.5 | 86.7 | 74.1 | • | 92.6 | |
| Ĭ | 4aii | Patient experience of primary care - GP out of hours services 2012-13 (Score out of 100) | 412 | 66.6 | 70.2 | 54.5 | • • | 82.8 | • |
| | 4aiii | Patient experience of primary care - NHS dental services 2012-13 (Score out of 100) | 1626 | 89.2 | 84.0 | 73.4 | ♦ • | 92.9 | • |
| | 4b | Patient experience of hospital care 2013-14 (Score out of 100) | | 74.9 | 76.9 | 67.1 | • | 87.0 | • |
| | 4ci | Friends and Family Test - A&E Oct-13 (Score between -100 and 100) | | 61.0 | | -11.0 | | 93.0 | _ |
| | 4cii | Friends and Family Test - Inpatients Oct-13 (Score between -100 and 100) | | 76.0 | 71.0 | 41.0 | • | 100.0 | • |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Source

Public Health outcomes framework

NHS outcomes framework

Adult social care outcomes framework

National average

Dudley LA peers

Worst ● ◆ Best

| Domain | Indicator | | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------|-----------|--|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 4.1 | Patient experience of outpatient services 2011 (Score out of 100) | | 77.4 | 79.5 | 71.0 | • | 88.5 | • |
| | 4.2 | Responsiveness to in-patients' personal needs 2012-13 (Score out of 100) | | 66.5 | 68.7 | 54.4 | • | 84.2 | • |
| | 4.3 | Patient experience of accident and emergency services 2012 (Score out of 100) | | 78.5 | 79.1 | 0.0 | • | 86.6 | • |
| | 4.4i | Access to GP services 2012-13 (%) | 3932 | 73.3 | 76.3 | 59.3 | •• | 86.8 | • |
| Se | 4.4ii | Access to NHS dental services 2012-13 (%) | 1751 | 95.9 | 93.0 | 83.0 | ♦ • | 97.4 | • |
| Healthy Services | 4.5 | Women's experience of maternity services (Score out of 100) | | | 78.9 | | | | • |
| ealthy (| 4.6 | Bereaved carers' views on the quality of care in the last 3 months of life (%) | | | 77.3 | | | | |
| Ĭ I | 4.9 | People's experience of integrated care (TBC) | | | | | | | |
| | 5a | Patient safety incidents reported (Rate per 100 admissions) | | | 715.4 | | | | |
| | 5b | Safety incidents involving severe harm or death (Rate per 100 admissions) | | | 4.8 | | | | |
| | 5c | Hospital deaths attributable to problems in care (TBC) | | | | | | | |
| | 5.1 | Incidence of healthcare-related venous thromboembolism (Rate per 100,000 population) | | | | | | | |



Domain Indicator



HEALTH AND WELLBEING PRIORITIES -OUTCOMES FRAMEWORKS

Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

National average Dudley LA peers Worst **♦ Best**

> 25th Pecentile 75th Eng Previous **England Range Eng Best** Worst

| SS | 5.2i | Incidence of healthcare associated infection: MRSA bacteraemia 2012-13 (Number) | | 2.0 | | 18.0 | | 0.0 | |
|------------------|---------|--|-------|-------|-------|-------|---------------|------|---|
| | 5.2ii | Incidence of healthcare associated infection: C. Difficile 2012-13 (Number) | | 106.0 | | 269.0 | | 12.0 | |
| | 5.3 | Incidence of newly-acquired category 2,3 and 4 pressure ulcers (%) | | | | | | | |
| | 5.4 | Incidence of medication errors causing serious harm (Rate per 100 admissions) | | | 0.3 | | | | • |
| | 3.3xiii | PPV vaccination coverage (over 65s) 2012-13 (%) | 42512 | 69.4 | 69.1 | 55.3 | * | 77.0 | • |
| Service | 3.3xiv | Flu vaccination coverage (over 65s) 2012-13 (%) | 44871 | 72.7 | 73.4 | 65.5 | (| 80.8 | • |
| Healthy Services | 3.3xv | Flu vaccination coverage (at risk individuals from age six months to under 65 years) 2012-13 (%) | 17120 | 53.0 | 51.3 | 44.2 | \omega | 68.8 | |
| Ĭ | 3.4 | People presenting with HIV at a late stage of infection 2010-12 (%) | 12 | 48.0 | 48.3 | 80.0 | ♦ • | 0.0 | |
| | 3.5i | Proportion of patients who successfully complete treatment for tuberculosis 2012 (%) | | 74.3 | 82.8 | 100.0 | | 22.6 | |
| | 4.12i | Crude rate of sight loss due to Age-related Macular Degeneration (AMD) in persons aged 65 years and over 2012/13 (Rate per 100,000 population) | 94 | 155.6 | 104.4 | 221.3 | • | 31.7 | |
| | 4.12ii | Crude rate of sight loss due to glaucoma in persons aged 40 years and over 2012/13 (Rate per 100,000 population) | 12 | 7.3 | 12.5 | 29.3 | | 2.8 | |
| | 4.12iv | Crude rate of sight loss certifications 2012/13 (Rate per 100,000 population) | 168 | 53.6 | 42.3 | 79.8 | • | 13.5 | • |

Local

Numerator

Local

Value

Eng

Avg





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

| Domain | Indicator | • | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|-----------|-----------|---|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 2.1 | Proportion of people feeling supported to manage their condition 2012-13 (%) | | 63.7 | 65.6 | 54.3 | • | 78.9 | • |
| | 1.8i | Gap in the employment rate between those with a long- term health condition and the overall employment rate 2012 (%) | | 4.7 | 7.1 | 21.7 | ♦ | -5.3 | • |
| | 1.8ii | Gap in the employment rate between those with a learning disability and the overall employment rate 2011-12 (%) | | 65.7 | 63.2 | 73.1 | • • | 40.2 | • |
| | 1G | Proportion of adults with a learning disability who live in their own home or with their family 2013-14 (%) | 595 | 74.8 | 74.8 | 47.7 | ♦ | 94.5 | • |
| S | 1E | Proportion of adults with learning disabilities in paid employment 2013-14 (%) | 55 | 7.0 | 6.8 | 0.9 | ♦ | 23.1 | |
| Services | 1A | Social care related quality of life 2013-14 (Score out of 24) | 590 | 19.4 | 19.0 | 17.7 | ♦ • | 20.5 | |
| Healthy (| 1B | The proportion of people who use services who have control over their daily life 2013-14 (%) | 625 | 77.5 | 76.7 | 64.4 | ⋄ • | 90.9 | |
| ヹ | 1Cii | Proportion of people using social care who receive self-directed support 2013-14 (%) | 4135 | 63.5 | 62.1 | 25.4 | * | 138.1 | • |
| | 1Cii | Proportion of people using social care who receive direct payments 2013-14 (%) | 1260 | 19.3 | 19.1 | 6.1 | • | 47.1 | • |
| | 1D | Carer-reported quality of life 2012-13 (Score out of 12) | 355 | 8.3 | 8.1 | 6.6 | | 9.8 | • |
| | 11 | Proportion of people who use services and their carers, who reported that they had as much social contact as they would like (%) | | | | | | | |
| | 2Ai | Permanent admissions to residential and nursing homes of younger people, per 100,000 population 2013-14 (Rate per 100,000 population) | 35 | 18.3 | 14.4 | 44.8 | • | 3.6 | |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average

Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

| Domain | Indicato | r | Local Numerator | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------|----------|---|--------------------|----------------|------------|--------------|---------------|----------|----------------|
| | 2Aii | Permanent admissions to residential and nursing homes of older people, per 100,000 population 2013-14 (Rate per 100,000 population) | 480 | 794.5 | 668.4 | 1277.4 | • | 207.5 | • |
| | 2Bi | Proportion of older people (65 and over) who were still at home 91 days after discharge from hospital into reablement/rehabilitation services 2013-14 (%) | 265 | 87.4 | 81.9 | 58.9 | ♦ • | 100.0 | • |
| | 2Bii | The proportion of older people aged 65 and over offered reablement services following discharge from hospital. 2013-14 (%) | 300 | 3.3 | 3.3 | 0.6 | • | 25.8 | • |
| | 2Ci | Delayed transfers of care from hospital per 100,000 population 2013-14 (Rate per 100,000 population) | 27 | 10.8 | 9.7 | 27.2 | © | 1.3 | • |
| S | 2Cii | Delayed transfers of care from hospital attributable to adult social care services per 100,000 population 2013-14 (Rate per 100,000 population) | 18 | 7.3 | 3.1 | 13.7 | • • | 0.3 | |
| Healthy Services | 2D | The outcomes of short-term services: sequel to service (%) | | | | | | | |
| althy (| 2E | The effectiveness of reablement services (TBC) | | | | | | | |
| Ĭ | 2F | Dementia - a measure of the effectiveness of post-diagnosis care in sustaining independence and improving quality of life (TBC) | | | | | | | |
| | 3A | Overall satisfaction of people who use services with their care and support 2013-14 (%) | | 63.8 | 64.9 | 45.5 | | 83.7 | |
| | 3B | Overall satisfaction of carers with social services 2012-13 (%) | 155 | 45.9 | 42.7 | 25.8 | • | 65.4 | • |
| | 3C | The proportion of carers who report that they have been included or consulted in discussion about the person they care for 2012-13 (%) | 215 | 74.5 | 72.9 | 55.0 | 0 | 100.0 | • |
| | 3D | The proportion of people who use services and carers who find it easy to find information about services 2013-14 (%) | | 76.7 | 74.7 | 65.0 | ♦ • | 86.4 | • |





Spine chart explanation:

Significantly better than England average

Not significantly different from England average
Significantly worse than England average

No significance can be calculated

Public Health outcomes framework

NHS outcomes framework

Source

Adult social care outcomes framework

National average

Dudley LA peers
Worst ● ♦ Best

| Domain | Indicator | Indicator | | Local Value | Eng Avg | Eng Worst | England Range | Eng Best | Previous value |
|------------------|-----------|---|--|----------------|------------|--------------|---------------|----------|----------------|
| Healthy Services | 3E | Improving people's experience of integrated care (TBC) | | | | | | | |
| | 4A | The proportion of people who use services who feel safe 2013-14 (%) | | 70.8 | 66.0 | 54.7 | ♦ • | 81.8 | • |
| | 4B | The proportion of people who use services who say that those services have made them feel safe and secure 2013-14 (%) | | 90.9 | 79.2 | 53.8 | \Q | 94.3 | |